



# Ridmus™ Alpha

## Rodenticide

syngenta®

**Active ingredient:** Alphachloralose 3.996% w/w (39.96 g/kg)

**AUTHORISATION NUMBER:** UK-2016-1033-0001

**UFI:** 72CW-6000-E00U-T17P

A READY-TO-USE (RB) PASTE (PA) BAIT. FOR USE ONLY BY PROFESSIONALS ONLY AS A RODENTICIDE.  
FOR USE ONLY IN THE CONTROL OF MICE, INDOORS.

**Use Biocides safely and sustainably. It is illegal to use this product for uses or in a manner other than that prescribed on this label. NOT for Amateur sale. Read product label before use.**

### Warning

Very toxic to aquatic life, with long lasting effects.

Avoid release to the environment.

Collect spillage.

Dispose of contents/container in accordance with local /regional/national requirements.



### DIRECTIONS FOR USE:

#### FOR MICE (*Mus musculus*, house mouse - adult and juveniles)

Apply 8g of paste in a tamper resistant bait box, using a caulking gun.

#### RECOMMENDED BAITING FREQUENCY

2 bait boxes (with 8g Ridmus Alpha each)/ m. Replenish bait when necessary, due to the products fast action one round of application may be sufficient.

The number of bait boxes used is dependent on the site of treatment and severity of the infestation. As a guide – Place bait boxes 2 to 3 metres apart (3 metres apart in case of low infestation to 2 metres apart in case of high mouse infestation). Check bait boxes frequently. Replace any eaten bait. Replace any mouldy or contaminated bait. If all bait is eaten increase number of bait boxes and / or visit frequency. Replace bait until consumption stops. Bait should be laid for periods at most 7-10 days. If control is not achieved survey site again to establish reasons and consider re-siting or increasing bait boxes.

## Contents: 300 g e

**Batch No./Expiry date:** See pack

**Label No.:** CLP20-068



Read label before use and follow instructions. The product must not be placed indiscriminately. A thorough survey of the infested area is essential particularly in secluded and sheltered places, to determine the extent of the infestation. Wash hands and directly exposed skin after handling the product and before eating, drinking or smoking. Place bait boxes where the rodents are seen in or near burrows and harbourages on runs and places where they find food or gnaw. The bait must be placed inside bait boxes out of reach of children, birds, pets and other non-target animals. Bait boxes must be clearly marked to show they contain rodenticides and must not be disturbed. Remove all baits after treatment. When control has been achieved consider control measures (plug holes, remove potential food etc.) to reduce likelihood of reinfestation. Optimal efficacy is achieved at low room temperatures i.e. preferably as low or lower than 16°C. Dead rodent bodies, remains of unused bait or any fragments of bait found away from the bait station must be collected during all control operations to minimise the risk of consumption and poisoning to children, companion animals and other non-target animals. Dead rodent bodies, remains of unused bait or any fragments of bait must be disposed of in accordance with local requirements. The bait must be placed inside bait boxes out of reach of children birds and other non-target animals. Dead rodent bodies, remains of unused bait or any fragments of bait must be disposed of in accordance with local requirements. Remove all remains of bait after treatment. The delay of action is between 24-48h.

**PRECAUTIONS:** Do not place bait boxes near water drainage systems where they can come into contact with water. To maximise bait take ensure that alternative food sources have been removed wherever possible. Do not use product where it can come into contact with food, drinking water, animal feeding stuffs, utensils that have contact with food. Do not clean bait boxes between applications (internal only). Rodents can be disease carriers (e.g. Leptospirosis).

**FIRST AID INSTRUCTIONS:** Excessive concentrations may cause nervous system depression, headache and weakness leading to unconsciousness. No antidote. Treat symptomatically. If swallowed seek medical advice immediately and show this container or label. Call a POISON CENTRE or doctor if you feel unwell.

**INSTRUCTIONS FOR SAFE DISPOSAL OF THE PRODUCT AND ITS PACKAGING:** Wear adequate protective gloves when disposing of rodents. In the UK poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site. For further information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>)

**CONDITIONS OF STORAGE AND SHELF LIFE OF PRODUCT UNDER NORMAL CONDITIONS OF STORAGE:** Store product only in closed original container in a dry cool place. Keep away from children and non-target animals. Keep away from food drink and animal feeding stuff. Shelf life: 12 months.

**APPROVAL HOLDER:** Rentokil Initial 1927 plc, Power Centre, Unit A1 and A2, Napier Way, Crawley, West Sussex RH10 9RA, UK.

**MANUFACTURED BY:** Rentokil Initial Supplies, Webber Road , Liverpool, L33 7SR UK

Product Advice Line: +44 (0) 151 548 5050 Emergency Line : +44 (0) 1342 833 022