



Talon[®] Soft XT

Rodenticide

FOR TRAINED PROFESSIONALS ONLY

Active ingredient: Brodifacoum 0.0025% w/w (0.025 g/kg)
Great Britain registration number: GB-2017-1115-3-0002
Northern Ireland registration number: NI-2017-1115-3-0002
Republic of Ireland registration number: IE/BPA 70662-3-002
Read product label and leaflet before use. Retain packaging for information.

APPLICATION DETAILS	
Rats (<i>Rattus norvegicus</i> , <i>Rattus rattus</i>) Indoor and outdoor use (around buildings)	Secure up to 200g of product in covered tamper resistant bait stations or covered bait points spaced 10m apart (5m apart in areas of high infestation) in areas where rats are active.
Mice (<i>Mus musculus</i>) Indoor and outdoor use (around buildings)	Secure up to 50g of product in covered tamper resistant bait stations or covered bait points spaced 10m apart (5m apart in high infestation areas) in areas where mice are active.
Regularly check bait consumption and replace consumed or spoilt bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings).	

DIRECTIONS FOR USE: Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it. Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation. Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve. The product should only be used as part of an integrated pest management (IPM) system including, amongst others, hygiene measures and, where possible, physical methods of control. The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc). Where possible bait stations must be fixed to the ground or other structures. Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened. Bait must be secured so that it cannot be dragged away from the bait station. Place the product out of reach of children, birds, pets and farm animals, and other non-target animals. Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these. Wear protective chemical resistant gloves during product handling phase (EN374). When using the product do not eat drink or smoke. Wash hands and directly exposed skin after using the product. The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice. If bait uptake is low relative to the size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation. If after a treatment period of 35 days, baits continued to be consumed and no decline in rodent activity can be observed, the likely cause must be determined. Where other elements have been excluded, it is likely that there are resistant rodents so consider the use of a non anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure. Use a spatula for post-application (cleaning/disposal) of bait.

RISK MITIGATION MEASURES: Where possible, prior to treatment, inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign in accordance with the code of best practice. **In Ireland:** When the product is being used in public areas and tamper resistant bait stations are not used, the areas treated must be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant, as well as indicating the first measures to be taken in case of poisoning, must be made available alongside the baits.

Do not use in areas where resistance to the active substance can be suspected. Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment. Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant. Do not wash the bait stations or utensils used in covered or protected bait points with water between applications. Dispose of dead rodents in accordance with local requirements.

In the UK: Poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site. For further information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements. Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Refer to the CRRU UK Code of Best Practice (or equivalent) for guidance. When this product is supplied to a user for the control of rodents, it shall only be supplied to a professional user holding certification demonstrating compliance with the UK rodenticide stewardship regime requirements.

In Ireland: This product should only be used in accordance with a code of best practice such as the CRRU Ireland Best Practice Requirements for Rodent Control and Safe Use of Rodenticides. Follow any additional instructions in that code of best practice. Trained professional users must register with DAFM as a 'Trained Professional - Pest Management Professional (PMU)' and must present their PMU No at point of sale to purchase this product for use only as specified on this label.

USE-SPECIFIC INSTRUCTIONS FOR USE AND RISK MITIGATION MEASURES	
INDOOR USE	Remove the remaining product at the end of the treatment period. Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign. Consider preventative control measures (e.g. plug holes, remove potential food and drink as far as possible) to improve product intake and reduce the likelihood of reinfestation. To reduce the risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice. In Ireland these recommendations are provided by the CRRU Ireland Best Practice Requirements for Rodent Control and Safe Use of Rodenticides. Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities. Do not use this product in pulsed baiting treatments. When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.
OUTDOOR (AROUND BUILDINGS)	Protect bait from atmospheric conditions. Place the baiting points in areas not liable to flooding. Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt. Follow any additional instructions provided by the relevant code of best practice. Remove the remaining product at the end of the treatment period. For outdoor use, baiting points must be covered and placed in strategic sites to minimise exposure to non target species. Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign in accordance with the code of best practice. Consider preventative control measures (e.g. plug holes, remove potential food and drink as far as possible) to improve product intake and reduce the likelihood of reinfestation. To reduce the risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice. In Ireland these recommendations are provided by the CRRU Ireland Best Practice Requirements for Rodent Control and Safe Use of Rodenticides. Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities. Do not use this product in pulsed baiting treatments. Do not apply this product directly in burrows. When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

OTHER INFORMATION: Because of their delayed mode of action, anticoagulant rodenticides take from 4 days to 10 days to be effective after consumption of bait.

Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing of them. This product contains a bittering agent and a dye.

Authorisation holder (Great Britain): Rentokil Initial 1927 plc, The Power Centre, Unit A1 & A2 Link 10, Napier Way, Crawley, RH10 9RA, UK.

Authorisation holder (Republic of Ireland and Northern Ireland): Rentokil Initial Limited, Hazel House, Millennium Park, Naas, Ireland.

UK marketing company: Syngenta UK Limited. CPC4, Capital Park, Fulbourn, Cambridge, CB21 5XE. Tel: (01223) 883400

Irish marketing company: Syngenta Ireland Ltd. Industrial Park, Cork Road, Waterford. Tel :051-377203

Product Advice Line: +44 (0) 151 548 5050
Emergency Line: +44 (0) 1342 833 022.

10 x 300 g

Batch No./Expiry date:
Label No.: CLP23-074b

Product names marked ® or ™, the ALLIANCE FRAME and the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company